

February 2024

ESIG has finalised its 2022 solvent volatile organic compounds (VOC) emission inventories. As there was a decline in sales due to the economic situation in Europe, we also see a decline in the calculated VOC emissions.

General trend and comparison (amounts in kilotons)

year	ESIG VOC	ESIG VOC emissions	EEA	EEA vs ESIG
	emissions	with ethanol	inventories*	difference
2008	2159	NA	3369	36%
2000	2133			3070
2009	1917	NA	3029	37%
2013	1775	2323	2706	16%
2015	1842	2366	2632	11%
2016	1628	2002	2617	31%
2017	1613	1980	2678	35%
2018	1813	2145	2644	23%
2019	1765	2120	2589	22%
2020	1538	2265	2751	21%
2021	1663	2209	2692	22%
2022	1470	1816		

^{*} based on reporting in 2023

Both ESIG and EEA data show relatively stabilized emission level in the later 2010s, the difference between the two datasets showing up to more than 30% divergences. In reporting of solvents since 2018 however, the difference decreased to slightly over 20%. In 2020, the ESIG VOC emissions decreased, which may be attributed to lower production/sales given the Covid-19 crisis. On the other hand, ethanol consumption increased given the use in hand sanitizer, which all in all led to an increase in VOC emissions overall compared to the previous years. In 2021 this increased use of ethanol was still found, albeit lower than in 2020. In 2022 there is a decline most likely due to the economic situation.

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium









February 2024

The ESIG solvents emission inventories have been conducted with the support of TNO.

ESIG Solvent VOC Emissions Numbers 2022 per country

Summary of results for aggregated country groups applied by ESIG 2022								
	TOTAL EMISSIONS BY COUNTRY OF PRODUCTION			Emissions				
Countries	Hydrocarbon+Oxygenated	IMPORT/EXPORT	Ethanol	NMVOC in				
Countries Austria	solvents (kiloton/country) 27.54	Corrections 10.11	emissions	kiloton/country				
			6.04 8.24	43.69				
Belgium + Luxembourg	122.48	-93.42		37.31				
Bulgaria + Romania	5.16	38.38	17.40	60.93				
Croatia	0.89	15.75	2.61	19.25				
Cyprus + Greece + Malta	13.16	15.73	8.09	36.98				
Czech Republic	11.30	26.14	7.07	44.50				
Denmark	16.46	-0.76	3.95	19.65				
Estonia + Latvia + Lithuania	5.43	21.73	4.04	31.19				
Finland	9.43	9.92	3.73	23.08				
France	181.73	-1.07	45.61	226.27				
Germany	429.68	-280.57	55.96	205.07				
Hungary	8.30	16.94	6.51	31.75				
Ireland	13.99	-0.86	3.40	16.53				
Italy	166.24	61.68	39.65	267.57				
Netherlands	160.02	-91.89	11.83	79.97				
Poland	41.65	55.54	25.31	122.51				
Portugal	12.16	37.84	6.96	56.96				
Slovakia	1.39	21.67	3.65	26.72				
Slovenia	10.72	-0.25	1.42	11.89				
Spain	98.63	75.63	31.89	206.15				
Sweden	29.22	11.31	7.03	47.55				
United Kingdom	104.14	50.45	46.09	200.68				
Total EU-27+UK	1469.73	0.00	346.48	1816.20				

Data on import/export are taken from J. Pearson for 2016 & 2017, based on adjusted 2015 import/export.

From 2018 onwards, import/export estimates are based on the import and export estimates from the year before which are adjusted for the increase/decrease in VOC emissions.

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium







February 2024

As from 2021, import and export estimates have been estimated for this year as well as revised for the years before. This does not affect EU28 totals but at country level adjustments have been made. The main goal of this update was to improve consistency in the methodologies for all years to the extent possible. Once the EEA data is available, we will add the comparison table.

ESIG Solvent VOC Inventories for the EU – a top-down approach¹

In summary, ESIG's top-down approach consists of three operations:

- 1. Collecting the solvent sales volumes in each EU country according to REACH end use sector
- 2. Applying the air emission factor for each use sector & calculating VOC emissions
- 3. Applying import/export corrections

1. All companies which are part of ESIG are invited to submit their solvent sales volumes by end-use sector for each European country (EU-27+ UK).

Notes:

- The sales numbers are about 90% of all oxygenated and hydrocarbon VOC solvents² manufactured and then sold in the European Union hence highly confidential. There are collated by Cefic, and carefully and confidentially analysed.
- No data from chlorinated/halogenated solvents are included anymore as all uses are in closed system, meaning no emission to air.
- Excluded from ESIG's portfolio is also any substance which main use is not as solvent such as Toluene, Benzene and Xylene DMF, NMP, THF Propylene Glycols Methanol
- The end-use sector are the same market sectors as defined for the Generic Exposure Scenario
 (GES) of the Registration, Evaluation and Authorisation of Chemicals Regulation (REACH). For the
 estimation of environmental exposure, ESIG has developed Specific Environmental
 Release Categories (SpERCs), which provide sector-specific release fractions, associated
 operational conditions and risk management measures.

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium





¹ For more information on the method used, former technical papers can be consulted: https://www.esig.org/wp-content/uploads/2018/03/201802_ESVOC_techncial-paper-solvent-VOC-emisisons-2015_final-1.pdf
https://www.esig.org/wp-content/uploads/2019/03/Atmospheric-Environment-John-VOC-article-201903.pdf

² Members are asked to report only solvent sales for VOC solvents according to the definition given in the Industrial Emissions Directive: 'volatile organic compound' means any organic compound (...), having at 293,15 K a vapour pressure of 0,01 kPa or more, or having a corresponding volatility under the particular conditions of use"



February 2024

- 2. The emissions factors (see table below) have been assessed by using the GES approach, and estimate the percentage of VOCs emitted into the air. The solvent VOC emissions per sector are then calculated by multiplying the solvent volumes by the emission factor for that sector. These emissions factors depend on two parameters: the final end-use of the solvent and the type of solvent. As each of these end-uses is associated with one or more generic exposure scenarios (GES) where the release-to-air percentages have been determined from the environmental safety assessment based on realistic data and/or assumptions for particular uses (2010). Not all solvents are VOC and not all VOC's are solvents. Their respective properties such as boiling point range, molecular weight make them VOC or not. For example, a C6 Hydrocarbon solvent (Hexane) will be a VOC whereas a heavier C14-C19 hydrocarbon solvent is not a VOC.
- 3. Due to the free flow of goods in the European Union, there are no direct data for import and export of solvents within the EU member states and so an estimate has to be made. Cefic (with as source Eurostat) has data for overall chemicals exports and imports for each country so that the net chemical transfer can be calculated for each Member State. By assuming this percentage of chemical transfer is directly proportional to percentage of solvent movements, an estimate of solvent import/export related to solvent downstream activities has been made for each country for 2015, which has been extrapolated to the later years. These calculations assume that there is no net import or export to or from the EU. These data show that Belgium, Germany and Netherlands are substantial exporters of solvents and downstream products (as such "exporting" VOC emissions) within the EU. The UK and France are effectively in balance as we assume is Ireland, and it is further assumed that there is no net import or export of solvents for these three countries. The remaining EU Member States are net importers of solvents, and although they export and import to one another, their total net imports are assumed to equal the exports from Belgium, Germany and the Netherlands. An apportion is made based on population data. For the 2021 ESIG emissions, the import and export assumptions have been reassessed, where especially for smaller countries adjustments were made.

Changes in 2013-2021 time series compared to 2016-2020 time series

By the end of 2022, the time series 2013 and 2015-2021 have been completed for the ESIG inventory. The results have been communicated to the inventory teams late 2022, early 2023.

The changes are the following:

- Extension of the time series, adding one year at the end (2021) and two years at the beginning (2013 and 2015).

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium









February 2024

- With the addition of the new years, import and export have been reassessed for the time series, which results in relatively small changes overall, however for some specific countries the differences can be larger.
- The calculation of ethanol use for hand sanitizers has been revised with new data available from the industrial branch organization (E-PURE) for the EU27+UK as a whole.
- The distribution of ethanol use over individual Member States has been revised. The old distribution was based on a Eurostat indicator for the use of personal care products, where some countries were missing and therefore received no emissions related to ethanol. Given the Covid-19 pandemic, ethanol has become a significantly more important source of emissions and the products were used by the general population. Therefore, a distribution to countries proportional to the total population in each year has been assumed in the new inventory. This causes quite some changes in each country, especially for 2020 where the impact of Covid-19 on ethanol-related VOC emissions is significant.
- Since ethanol emissions have been included in NFR category 2D3a, also the per capita emission factors for this specific source category have been revised in the new inventory, and in some cases, these are significantly different.

Comparison of the results

Because of the changes outlined above, the emission estimates for historical years (2016-2020) in the latest version may be different from the previous ESIG inventory. The changes are relatively larger for smaller countries as are the uncertainties, which relates to the difficulties in assessing the transport of solvents or solvent containing products between EU Member States prior to the use of them. At the overall EU27+UK level the total VOC emissions are 1-2% smaller in each of the years 2016-2020. At Member State level however, the differences may be significant, ranging from -68% (Latvia) to +31% (France) compared to the previous inventory. It should be stressed however that the new estimates are thought to represent better reality.

When looking at category 2D3a (domestic solvent use) only, similar observations can be made. Compared to the previous version of the ESIG inventory, the EU27+UK total difference for each of years 2016-2020 is maximum 2%. At the level of individual countries however, the differences are significantly larger, for 2020 ranging from -83% (Malta) to +75% (France). The main factor in this difference, for both 2D3a and the total VOC emissions, is the change made in the distribution of ethanol-related VOC emissions over individual countries as described above (emissions from ethanol are 100% allocated to category 2D3a).

The table below provides per capita emission factors for all sectors and for category 2D3a only derived from the ESIG VOC inventory results, for the current latest version (2013-2021) compared to the previous version (2016-2020). All numbers refer to the year 2020 and represent the ESIG VOC inventory including ethanol.

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium











February 2024

Per capita emissions (kg/person) derived for the year 2020 for all sectors and for NFR category 2D3a only

	All se	ctors	2D3a only			
Country	2016-2020	2013-2021	2016-2020	2013-2021		
AT	6.17	6.06	1.55	2.31		
BE	2.35	3.51	0.69	1.69		
BG	2.05	2.89	1.38	1.78		
HR	5.04	5.80	2.78	1.91		
CY	1.49	2.97	1.19	1.71		
CZ	5.50	5.16	1.10	2.16		
DK	3.58	4.23	1.83	2.73		
EE	2.72	5.28	0.94	2.65		
FI	6.51	5.16	1.89	2.43		
FR	5.32	4.05	3.66	2.09		
DE	2.77	3.34	0.94	1.76		
GR	3.13	3.82	1.58	1.82		
HU	3.08	3.90	0.80	1.85		
IE	3.97	4.09	0.47	1.73		
IT	7.32	5.76	2.96	1.94		
LV	2.05	6.31	1.04	2.38		
LT	1.99	5.78	0.47	2.22		
LU	2.22	4.25	1.09	1.80		
MT	1.64	3.47	0.31	1.81		
NL	4.54	5.84	0.80	2.19		
PL	3.88	4.15	2.09	1.84		
PT	1.80	5.25	0.60	2.32		
RO	2.41	3.20	0.61	1.58		
SK	2.19	5.93	1.32	4.14		
SI	5.11	7.02	1.33	1.87		
ES	5.75	5.29	3.09	2.46		
SE	4.69	4.89	1.44	2.02		
UK	3.34	3.86	2.00	2.00		
Average	4.31	4.40	2.00	2.02		

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium









February 2024

How to access and use ESIG data?

The ESIG inventories are an accepted Tier 2a method according to the EMEP/EEA air pollutant emission inventory guidebook, however, the top-down approach using REACH end uses and confidentiality considerations make it difficult to use the data when establishing the country inventories.

Due to the high confidentiality of sales data Cefic statistics impose certain rules. When there are fewer than 3 original entries there is no show of data for a category. In few cases countries are grouped together to allow to display at least some numbers. Here again population numbers could serve to split between countries to get some results.

Thanks to efforts done by the UK, a table is added as an annex to the EMEP/EEA air pollutant emission inventory guidebook 2019, chapter 2.D.3.a.Domestic solvent use mapping the NFR categories against Reach end use sectors showing where and when ESIG data can be used (see table below). For instance, for INDUSTRIAL CLEANING: 100% is attributed to 2D3e, so in case of no own activity data the ESIG number can be used instead.

Since ESIG data are based on real data from EU solvents manufacturers, the ESIG emission inventories remain a valuable source of information and can also be used for comparisons.

Member States can obtain more information upon request for full details per REACH category provided data is not confidential. Additionally, based on the mapping, ESIG can per member state, and provided no confidentiality claims apply, provide a reattribution to the NFR categories.

REACH end use categories & Emission Factors

	End-use	Release to air	Remarks
1	Agrochemical uses	100%	
2	Blowing Agents	100%	assumed to be completely released to the atmosphere
3	De-Icing	100%	and utilize solvents that are VOC
4	Binder and Release Agents	100%	
5	Industrial Cleaning	70%	Cleaning agents used industrially are mostly handled in a closed system and are partially released to atmosphere
6	Professional & Consumer Cleaning	50%	There are two types of products in this category: dry cleaning agents used by professionals that have very low release percentage and the other cleaning agents used by

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium









February 2024

		1	
			consumers that are completely released to atmosphere. Therefore, an average 50% emission factor is applied
7	Industrial, Professional and Consumer Coatings	75%	The emission factor of the entire coatings industry is a combination of consumer and professional paint for the decorative market which has an emission factor of 100% and the industrial paints where the solvents are mostly regenerated resulting in an emission factor assumed to be 10%. Therefore, a conservative emissions factor of 75% is applied.
8	Functional Solvents	10%	include solvents used in chemical processes including intermediates, polymerization and extraction resulting in a low emission factor of 10%.
9	Metal working/Rolling Oils/ Lubricant uses	0%	taken over from ATIEL, the technical Association of the European Lubricants Industry
10	Oil field chemicals-Drilling- Mining-Extraction	0%	Solvents used are mostly heavy and non-VOC
11	Polymers Processing (incl. rubber-tyre production)	10%	Value from ESIG GES/SpERCS
12	Road and Construction	95%	Value from ESIG GES/SpERCS
13	Use as Fuel/Combustion	0.25%	Combustion solvents are burnt, generating water and carbon dioxide and therefore do not produce any VOC in the atmosphere. A conservative emission factor of 0.25% has been applied to take into account possible leaks.
14	Water Treatment	5%	Value from ESIG GES/SpERCS
15	Other Consumer uses (household, aerosols, cosmetics)	90%	Solvents used by consumer in household and aerosol applications are completely released to the atmosphere. Solvents used in cosmetic applications. are heavy and non-VOC products. Therefore, an estimated 90% emission factor has been applied.
16	Pharmaceuticals Manufacturing	30%	Value from ESIG GES/SpERCS

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium











February 2024

Table annexed to the EMEP/EEA air pollutant emission inventory guidebook 2019, chapter 2.D.3.a.Domestic solvent use

REACH end-use sector	2D3a	2D3b	2D3c	2D3d	2D3e	2D3f	2D3g	2D3h	2D3i
Agrochemical uses	100 %								
Blowing agents									100 %
De-icing	50 %								50 %
Binder and release agents									100 %
Cleaning industrial and leather									
treatment					100 %				
Cleaning — professional									
consumer	100 %								
Coatings — industrial and									
adhesives, inks				80 %				15 %	5 %
Coatings —									
professional/consumer and									
thinners, paint industry	30 %			70 %					
Functional solvents (including									
solvents used in chemical									
processes, e.g. process aids,									
intermediates, extraction,									
dewaxing agents)							100 %		
Metal working/rolling									
oils/lubricant uses									100 %
Oil field chemicals, drilling,									
mining, extraction									100 %
Polymers processing (including									
rubber-tyre production) and									
industrial resins, synthetic									
rubber, process							100 %		
Road and construction		100 %							
Use as fuel/combustion and									
fuel additives									100 %
Water treatment									100 %
Other consumer uses									
(household, aerosols,									
cosmetics)	100 %								
Pharmaceuticals									
manufacturing							100 %		
Others — please specify below									100 %

ESIG (European Solvents Industry Group) Rue Belliard 40 b.15 B-1040 Brussels Belgium





